

Table 1. Estimated numbers of diagnoses of HIV/AIDS, by year of diagnosis and selected characteristics of persons, 1999–2002—30 areas with confidential name-based HIV infection reporting

	Year of diagnosis				Cumulative through 2002 ^a
	1999	2000	2001	2002	
Age at diagnosis (yrs)					
<13	187	163	206	162	3,219
13—14	28	31	33	30	391
15—24	2,646	2,803	2,926	2,926	40,896
25—34	7,817	7,386	7,221	7,338	115,468
35—44	9,115	9,289	9,119	9,450	100,294
45—54	3,887	4,212	4,408	4,675	37,763
55—64	1,112	1,250	1,303	1,450	10,647
≥65	382	386	427	432	3,455
Race/ethnicity					
White, not Hispanic	7,718	7,985	8,040	8,347	115,134
Black, not Hispanic	14,398	14,129	14,090	14,398	162,950
Hispanic	2,631	2,920	3,012	3,321	29,315
Asian/Pacific Islander	123	150	140	149	1,262
American Indian/Alaska Native	153	147	142	168	1,795
Exposure category					
Male adult or adolescent					
Male-to-male sexual contact	9,995	10,519	10,974	11,701	134,357
Injection drug use	3,029	2,940	2,715	2,757	43,426
Male-to-male sexual contact and injection drug use	1,066	962	948	943	17,758
Heterosexual contact	3,237	3,252	3,191	3,234	30,337
Other ^b	166	167	167	164	2,694
Subtotal	17,493	17,842	17,995	18,798	228,572
Female adult or adolescent					
Injection drug use	1,651	1,590	1,481	1,418	22,166
Heterosexual contact	5,708	5,777	5,826	5,949	56,492
Other ^b	132	149	136	136	1,678
Subtotal	7,491	7,516	7,442	7,503	80,336
Child(<13 yrs)					
Perinatal	174	151	192	144	2,754
Other ^c	14	12	14	18	465
Subtotal	187	163	206	162	3,219
Total^d	25,174	25,522	25,643	26,464	312,133

Note. These numbers do not represent actual cases in persons with a diagnosis of HIV/AIDS. Rather, these numbers are point estimates of cases diagnosed that have been adjusted for reporting delays and for redistribution of cases in persons initially reported without an identified risk. The estimates have not been adjusted for incomplete reporting.

Data include persons with a diagnosis of HIV infection. This includes persons with a diagnosis of HIV infection only, a diagnosis of HIV infection and a later AIDS diagnosis, and concurrent diagnoses of HIV infection and AIDS.

Since 1998, the following 30 areas have had laws or regulations requiring confidential name-based HIV infection reporting: Alabama, Arizona, Arkansas, Colorado, Florida, Idaho, Indiana, Iowa, Louisiana, Michigan, Minnesota, Mississippi, Missouri, Nebraska, Nevada, New Jersey, New Mexico, North Carolina, North Dakota, Ohio, Oklahoma, South Carolina, South Dakota, Tennessee, Utah, Virginia, West Virginia, Wisconsin, Wyoming, and the U.S. Virgin Islands. Since July 1997, Florida has had confidential name-based HIV infection reporting only for new diagnoses.

^a Includes persons with a diagnosis of HIV infection, from the beginning of the epidemic through 2002.

^b Includes hemophilia, blood transfusion, perinatal, and risk not reported or not identified.

^c Includes hemophilia, blood transfusion, and risk not reported or not identified.

^d Includes persons of unknown or multiple race and of unknown sex. Cumulative total includes 1677 persons of unknown or multiple race and 6 persons of unknown sex. Because column totals were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.